

A NOTE ON QUOTIENTS OF WORD HYPERBOLIC GROUPS WITH PROPERTY (T)

YVES DE CORNULIER

ABSTRACT. Every discrete group with Kazhdan's Property (T) is a quotient of a torsion-free, word hyperbolic group with Property (T).

All groups in the paper are discrete and countable. Recall [HV] that a group G has Property (T) if every isometric action of G on an affine Hilbert space has a fixed point (or, equivalently, has bounded orbits).

As an immediate consequence of the definition, Property (T) is inherited by quotients and by extensions.

Shalom [Sha, Theorem 6.7] has proved the following interesting result about Property (T).

Theorem 1 (Shalom, 2000). *For every group G with Property (T), there exists a finitely presented group G_0 with Property (T) which maps onto G .*

In other words, this means that, given a finite generating subset for G , only finitely many relations suffice to imply Property (T). This can be interpreted in the topology of marked groups [Cha] as: Property (T) is an open property. See [Gr, 3.8] for a generalization to other fixed point properties.

A word hyperbolic group is a finitely generated group whose Cayley graph satisfies a certain condition, introduced by Gromov, meaning that, at large scale, it is negatively curved. We refer to [GH] for a precise definition that we do not need here. We only mention here that word hyperbolic groups are necessarily finitely presented, that word hyperbolicity is a fundamental notion in combinatorial group theory as in geometric topology. Word hyperbolic groups are groups with “many” quotients, and thus can be considered as a generalization of free groups.

It was asked [Wo, Question 16] whether every group with Property (T) is quotient of a group with Property (T) with finiteness conditions stronger than finite presentation. We give an answer here by showing that we can impose word hyperbolicity.

Proposition 2. *For every group G with Property (T), there exists a torsion-free word hyperbolic group G_0 with Property (T) which maps onto G .*

Note that Proposition 2 contains Theorem 1 as a corollary; however it is proved by combining Theorem 1 with the following remarkable result of Ollivier and Wise [OW]. Since it involves some technical definitions, we do not quote it in full generality.

Theorem 3 (Ollivier and Wise, 2005). *To every finitely presented group Q , we can associate a short exact sequence $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ such that*

- (1) G is torsion-free, word hyperbolic,
- (2) N is 2-generated and has property (T).

Corollary 4. *For every finitely presented group Q with Property (T), there exists a torsion-free word-hyperbolic group G with Property (T) mapping onto Q with finitely generated kernel.*

Proof: Apply Theorem 3 to Q , so that G lies in an extension $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$, where N has Property (T) and Q has Property (T). Since Q has Property (T) and since Property (T) is stable under extensions, G also has Property (T). ■

Remark 5. Corollary 4 answers a question at the end of [OW].

Date: February 1, 2008.

2000 Mathematics Subject Classification. Primary 20F65; Secondary 20F67.

Proof of Proposition 2: Let G be a group with Property (T). By Theorem 1, there exists a finitely presented group Q with Property (T) mapping onto G , and by Corollary 4, there exists a torsion-free word hyperbolic group G_0 with Property (T) mapping onto Q , so that G_0 maps onto G . ■

Question 6. 1) In Theorem 3, can G be chosen, in addition, residually finite? In [Wis], a similar result is proved, G being torsion-free, word hyperbolic, residually finite, and N finitely generated, but never having Property (T).

2) Let G be a word hyperbolic group (maybe torsion-free), and H a quotient of G generated by r elements. Does there exist an intermediate quotient which is both word hyperbolic and generated by r elements? (The analog statement with “word hyperbolic” replaced by “finitely presented” is immediate.) The motivation is that, in Proposition 2, we would like to have G_0 generated by no more elements than G . Theorem 3 only tells us that if G is r -generated, then G_0 can be chosen $(r + 2)$ -generated.

Question 7. Following [Ser], a group has Property (FA) if every isometric action on a simplicial tree has a fixed point. Is it true that every group with Property (FA) is a quotient of a finitely presented group with Property (FA)? The author imprudently claimed that this is true in an earlier version of this paper, but the proof contained a mistake.

REFERENCES

- [Cha] Christophe CHAMPETIER. *L'espace des groupes de type fini*, Topology **39**(4) 657-680, 2000.
- [GH] Étienne GHYS, Pierre DE LA HARPE. “Sur les groupes hyperboliques, d’après Mikhael Gromov”. Progress in Mathematics, 83. Birkhäuser, 1990.
- [Gr] Misha GROMOV. *Random Walk in Random Groups*. Geom. Funct. Anal. **13**(1), 73-146, 2003.
- [HV] Pierre DE LA HARPE, Alain VALETTE. “La propriété (T) de Kazhdan pour les groupes localement compacts”, Astérisque **175**, SMF, 1989.
- [OW] Yann OLLIVIER, Daniel WISE. *Kazhdan groups with infinite outer automorphism group*. To appear in Trans. A.M.S., 2005.
- [Ser] Jean-Pierre SERRE. “Arbres, amalgames, SL_2 ”. Astérisque **46**, SMF, 1977.
- [Sha] Yehuda SHALOM. *Rigidity of commensurators and irreducible lattices*. Inv. Math. **141**(1):1-54, 2000.
- [Wis] Daniel T. WISE. *A residually finite version of Rips’s construction*. Bull. London Math. Soc. **35**(1): 23-29, 2003.
- [Wo] Report of the workshop *Geometrization of Kazhdan’s Property (T)* (organizers: B. Bekka, P. de la Harpe, A. Valette; 2001). Unpublished; currently available at <http://www.mfo.de/cgi-bin/tagungsdb?type=21&tnr=0128a>.

Yves de Cornulier

École Polytechnique Fédérale de Lausanne (EPFL)

Institut de Géométrie, Algèbre et Topologie (IGAT)

CH-1015 Lausanne, Switzerland

E-mail: decornul@clipper.ens.fr